



COMPOSITION

Hexitiazox: 25.13% w/v (24.59% w/w).
Concentrated suspension (Cs).

USES

A non-systemic acaricide with contact and ingestion activity, good translaminar activity and a long-lasting residual effect. It has ovicidal, larvicidal and nymphicidal activity against numerous species of mites.

AUTHORISED USES

- **CITRUS FRUITS:** Tetranychid mites (*Tetranychidae*).
- **CUCURBITS WITH INEDIBLE SKIN:** Common spider mite (*Tetranychus spp.*)
- **POME FRUIT TREES:** Common spider mite (*Tetranychus spp.*)
- **TABLE GRAPES AND WINE GRAPES:** Common spider mite (*Tetranychus spp.*).

AUTHORISED MINOR USES (under national procedures)

- **APRICOT, PERSIMMON, CHERRY, PLUM, BLACKTHORN and NUT TREES:** Common spider mite (*Tetranychus spp.*).

DOSAGE AND METHOD OF USE

Outdoor insecticide/acaricide treatment by mechanical and manual foliar spraying.

Apply by standard foliar spraying at the onset of infestation, ensuring thorough coverage of all vegetation, particularly the undersides of the leaves. Treat eggs and larvae before they reach the adult stage. To control mites at all stages of their life cycle, it should be used in combination with an adulticide.

In **APRICOT, CHERRY, PLUM, BLACKTHORN AND NUT TREES** (minor uses), apply 160 to 300 mL of product per hectare with a spray mixture volume of 800 to 1,000 L/ha. Apply a single treatment during BBCH 00–69.

In **PERSIMMON** (minor use) apply 160 to 200 mL of product per hectare with a spray mixture volume of 800 to 1,000 L/ha. Apply a single treatment before fruit set, up to the end of flowering (BBCH 00–69).

In **CITRUS FRUITS** apply 40 to 150 mL of product per hectare with a spray mixture volume of 1,000 to 2,500 L/ha. Apply a single treatment during BBCH 00–97.

In **CUCURBITS WITH INEDIBLE SKIN** apply 60 to 150 mL of product per hectare with a spray mixture volume of 300 to 500 L/ha. Apply a single treatment during BBCH 10–89.

In **POME FRUIT TREES** apply 160 to 300 mL of product per hectare with a spray mixture volume of 800 to 1,000 L/ha. Apply a single treatment during BBCH 00–97.

In **TABLE GRAPES AND WINE GRAPES** Apply 100 to 300 mL of product per hectare with a spray mixture volume of 500 to 1,000 L/ha. Apply a single treatment during BBCH 00–97.

PRE-HARVEST INTERVAL

- **Cucurbits with inedible skin:** 3 days.
- **Citrus fruits:** 14 days.
- **Table grapes and wine grapes:** 21 days.
- **Persimmons and pome fruit trees:** 28 days.
- **Apricot, cherry, plum, blackthorn and stone fruit trees:** No pre-harvest interval is required.

PRECAUTIONS

Do not mix with pyrethroid insecticides.

HAZARD CLASSIFICATION



REGISTRY NUMBER

ES-01114