

CROP	PEST/EFFECT	DOSAGE
Tobacco	Aphids, <i>Aphididae</i> Lepidoptera Thrips	30-50 mL/hL
Wheat	Bugs, Heteroptera	50 mL/hL
	Aphids, <i>Aphididae</i> Thrips	30 mL/hL
Plant species (grown outdoors and in greenhouses, exclusively for use in nurseries)	Aphids, <i>Aphididae</i> Bugs, Heteroptera Weevils, <i>Curculionidae</i> Noctuids, <i>Noctuidae</i>	50 mL/hL
Ornamental shrubs and small trees (grown outdoors and in greenhouses)	Aphids, <i>Aphididae</i> Weevils, <i>Curculionidae</i> Noctuids, <i>Noctuidae</i>	50 mL/hL
Herbaceous ornamental plants (grown outdoors and in greenhouses)	Aphids, <i>Aphididae</i> Weevils, <i>Curculionidae</i> Spittlebugs, <i>Philaenus spumarius</i> Noctuids, <i>Noctuidae</i>	50 mL/hL
	Aphids, <i>Aphididae</i> Weevils, <i>Curculionidae</i> Whiteflies, <i>Aleyrodidae</i> Thrips	50 mL/hL
Herbaceous ornamental plants for cut flowers (grown outdoors and in greenhouses)	Cacoecia, the carnation caterpillar, <i>Cacoecimorpha pronubana</i> Beetles, <i>Scarabaeidae</i>	50-100 mL/hL
	Aphids, <i>Aphididae</i> Vine leaf beetle, <i>Byctiscus betulae</i> Spittlebug, <i>Philaenus spumarius</i> Noctuids, <i>Noctuidae</i> Grapevine moth, <i>Sparganothis pilleriana</i> Grape berry moth, <i>Lobesia botrana</i>	40-60 mL/hL
Carrots	Aphids, <i>Aphididae</i> Bugs, Heteroptera Whiteflies, <i>Aleyrodidae</i> Carrot fly, <i>Chamaepsila rosae</i>	50 mL/hL
	Lepidoptera Noctuids, <i>Noctuidae</i> Thrips	30 mL/hL

Apply by standard spraying outdoors using a backpack sprayer, a tractor-mounted sprayer for tall crops and a tractor-mounted sprayer for low crops, and using a lance or spray gun in greenhouses (only for cut flowers, ornamental plants and nurseries). Use 1,000 litres of mixture per hectare. Apply a maximum of two treatments per season at 14-day intervals, except for vines, where only one treatment per season should be carried out.

PRE-HARVEST INTERVAL

Lucerne and Cotton Plant: 15 days.

Chicory (leaves), Celery, Lamb's lettuce, Aubergine, Watercress, Shepherd's purse, Young shoots, including Brassica species, Wintercress, Barley, Onion, Campion, Dandelion, Strawberry, Stone fruit trees, Green peas, Fodder broad beans, Field beans, Green broad beans, Green beans, Maize, Chinese mustard, Cucumber, Pepper, Rocket, Tobacco, Tomato, Grapevine, Carrot: 3 days.

Pome fruit trees: 3 days (Pear: 7 days)

Almond, Rapeseed and Wheat: 30 days.

Olive, Potato, Leek and Cabbage: 7 days.

Ornamental shrubs and small trees, Poplars and aspens, Plant species, Herbaceous ornamentals: No pre-harvest interval is required.

HAZARD CLASSIFICATION



REGISTRY NUMBER

25,700