


FERTILUQ K-30



COMPOSITION

30% w/w Water-soluble Potassium Oxide (K_2O).

USES

FERTILUQ K-30 supplies potassium and corrects potassium deficiencies in all types of crops.

It is recommended that FERTILUQ K-30 be applied at times when the need for this element is greatest, such as during the fruit-filling stage (it increases fruit size, improves turgidity and raises sugar levels in the fruit).

It can be used on vegetable crops, ornamental plants, fruit trees, citrus trees, vines, olive trees and cereals.

DOSAGE AND METHOD OF USE

It is applied as a foliar spray at a rate of 150–300 cm^3/hL , depending on the severity of the deficiency and the timing of application. Depending on the circumstances, 2 or 3 applications may be necessary; it is recommended that these be started around 60 days before harvest or fruit ripening, or when a potassium deficiency is detected in the crop.

FERTILUQ K-30 can also be applied directly via fertigation as a supplement to the most suitable NPK fertiliser for each crop; it is therefore recommended to apply 10 to 20 L/ha, spread over two or three applications. It is recommended that the product be applied at a maximum pH of 7, as this helps prevent the drippers from becoming blocked and facilitates absorption by the plant.

PRECAUTIONS FOR USE

It is advisable to avoid applying the product during periods of high temperatures and strong sunlight, as this may cause the tips of the leaves to burn.

For further information, please contact our Agricultural Technical Department.

HAZARD CLASSIFICATION

