



COMPOSITION

35% w/w total phosphorus pentoxide (P_2O_5).

35% w/w total water-soluble Phosphorus Pentoxide (P_2O_5) total.

35% w/w phosphorus pentoxide (P_2O_5) soluble in neutral ammonium citrate.

35% w/w Water-soluble Potassium Oxide (K_2O).

Micronutrients

0.025% w/w Boron (B), water-soluble as boric acid.

0.01% w/w Water-soluble Copper (Cu) chelated with EDTA.

0.04% w/w Water-soluble Iron (Fe) chelated with EDTA.

0.07% w/w Water-soluble Manganese (Mn) chelated with EDTA.

0.003% w/w Molybdenum (Mo), water-soluble as sodium salt.

0.015% w/w Water-soluble Zinc (Zn) chelated with EDTA.

The pH range within which good stability of the chelated fraction is guaranteed is: 4–9.

USES

FERTILUQ S 0-35-35 is a soluble solid foliar fertiliser with a high phosphorus and potassium content, as well as chelated micronutrients. Formulated using high-purity raw materials with a high degree of solubility, FERTILUQ S 0-35-35 is characterised by its rapid and complete uptake by the plant.

It is particularly recommended for use before and after flowering to improve flowering and fruit set, and subsequently to enhance fruit size and ripening.

As a foliar fertiliser, it complements a balanced soil fertilisation regime. It can be used on all types of crops, particularly those where the quality and colour of the fruit are important, such as fruit trees, citrus trees, vines, olive trees, potatoes, beetroot and fruiting vegetables.

DOSAGE AND METHOD OF USE

Apply FERTILUQ S 0-35-35 as a foliar spray at a rate of 150–250 g/hL.

For tomatoes, apply 1–2 treatments after fruit set on each flower cluster at the recommended dose.

PRECAUTIONS FOR USE

Avoid applying the treatment during the hottest part of the day, as rapid evaporation of the water hinders its absorption by the leaves.

It can be used in combination with most plant protection products. Do not mix with oils or strongly alkaline products.

For further information, please contact our Agricultural Technical Department.

HAZARD CLASSIFICATION

