



COMPOSITION

6% w/w Water-soluble Copper (Cu) in gluconate form.

USES

A copper deficiency corrector for all types of crops. Its complexed formulation is the most suitable form for correcting copper deficiencies in any application.

DOSAGE AND METHOD OF USE

Leaf application: Apply as a standard foliar spray at a rate of 150–300 cm³ per hectolitre of water. Apply a sufficient amount of spray to ensure that all parts of the plant are completely covered. Depending on the severity of the deficiency, 2 to 3 spray applications are recommended, spaced 20 days apart, at the start of the growing season and before summer growth in citrus trees.

Application by fertigation: Apply with irrigation water at a rate of 1–3 L/ha per application. Depending on the severity of the deficiency, 4 or 6 treatments will be required, spread out over the growing season for fruit trees and citrus crops. Under no circumstances should the concentration exceed 0.05%.

PRECAUTIONS FOR USE

Avoid carrying out the treatment during the hottest part of the day.

Do not mix with oils (or substances containing oils).

Take precautions due to the phytotoxicity associated with copper, particularly in humid areas and with certain varieties of fruit trees and other crops.

Do not mix with strongly alkaline products, as the rapid evaporation of water hinders foliar uptake.

For further information, please contact our Agricultural Technical Department.

HAZARD CLASSIFICATION

