



COMPOSITION

5% w/w Water-soluble Iron (Fe), chelated with HEEDTA

The pH range within which good stability of the chelated fraction is guaranteed is: 4–9.

USES

A liquid iron supplement characterised by its high solubility and rapid uptake by all types of fruit trees, citrus trees, hazelnut trees, subtropical crops, vines and vine trellises, vegetables, ornamental plants and extensive crops.

DOSAGE AND METHOD OF USE

LUQSAFER L-65 is used diluted with water in any of the following systems: fertigation, hydroponics, via a localised soil applicator, injected into the soil or via trenches dug around the tree.

LUQSAFER L-65 should be used at the times and in the doses specified in the table.

| CROPS | | SOIL APPLICATION RATE | DRIP IRRIGATION |
|---|---|---|--|
| Fruit trees, citrus trees, hazelnut trees and subtropical trees | Seedlings | 5–25 cm ³ per tree | 3–15 cm ³ per tree |
| | Small trees | 25–50 cm ³ per tree | 15–30 cm ³ per tree |
| | Medium-sized trees | 50–100 cm ³ per tree | 30–65 cm ³ per tree |
| | Fully grown trees or trees showing severe chlorosis | 100–150 cm ³ per tree | 65–100 cm ³ per tree |
| | Period | From the start of budding and throughout the growing cycle | |
| Vines | Young vines | 5–15 cm ³ per vine | 3–10 cm ³ per vine |
| | Bearing vines | 15–25 cm ³ per vine | 10–15 cm ³ per vine |
| | Trellised vines | 25–50 cm ³ per vine | 15–30 cm ³ per vine |
| | Period | Before bud break and during the growing season | |
| Nurseries | - | 2–5 cm ³ per tree or 5–10 cm ³ per m ² | 1–3 cm ³ per tree or 3–6 cm ³ per m ² |
| | Period | Throughout the growing season | |
| Horticultural crops and ornamentals | - | 3–10 cm ³ per m ² | 2–6 cm ³ per m ² |
| | Period | After planting | |
| Extensive crops | - | 4–6 L/ha | - |

* It is recommended that these total doses be divided into 2 or 3 applications.

PRECAUTIONS FOR USE

Wash thoroughly with soap and water after handling.

Although it can be mixed with any type of fertiliser, it is advisable to check that the properties of both products are compatible.

To help the product penetrate down to the root level, we recommend watering the area after application.

For further information, please contact our Agricultural Technical Department.

