



COMPOSITION

Pyriproxyfen 10% w/v (100 g/L)
Emulsifiable concentrate (EC).

It contains petroleum naphtha (CAS 64742-47-8) and 2-ethylhexyl S-lactate (CAS 186817-80-1).

AUTHORISED USES

- **CITRUS FRUITS:** Caparreta, White louse, Grey louse, California red louse and Serpeta.
- **STONE FRUIT TREES:** San José scale.
- **POME FRUIT TREES:** San José scale.
- **TOMATO:** Whitefly.

AUTHORISED MINOR USES

- **COTTON PLANT:** Whitefly.
- **ALMOND TREE:** Mealybugs and San José scale.
- **HAZEL, CHESTNUT, WALNUT, PECAN, PECAN NUT and PISTACHIO:** Mealybugs,
- **AUBERGINE:** Whitefly.
- **PERSIMMON:** Mealybugs.

Apply as a standard foliar spray during the early larval stages of the target insects, preferably in the first generation.

- **COTTON** (minor use): apply a single treatment per growing season at a rate of 0.5 to 0.75 L/ha with a spray mixture volume of 1,000 L/ha, before the bolls open.
- **ALMOND, HAZELNUT, CHESTNUT, WALNUT, PECAN, PECAN NUT and PISTACHIO** (minor uses): apply a single treatment before flowering at a dose of 40–50 mL per 100 litres of water (0.04–0.05%), with a spray mixture volume of 700–1,000 L/ha, without exceeding 0.5 litres of product per hectare.
- **AUBERGINE** (minor use): only in greenhouse cultivation. Apply up to 2 treatments per growing season at a rate of 50–75 mL per 100 litres of water (0.05–0.075%), with a spray mixture volume of 1,000–1,500 L/ha, without exceeding 0.8 L/ha per application.
- **PERSIMMON** (minor use): apply a single treatment before flowering at a dose of 40–50 mL per 100 litres of water (0.04–0.05%), with a spray mixture volume of 700–1,000 L/ha, without exceeding 0.5 litres of product per hectare.
- **CITRUS FRUITS:** apply a single treatment per growing season at a dose of 50–75 mL per 100 litres of water (0.05–0.075%), with a spray mixture volume of 1,000–1,500 L/ha, without exceeding 1.125 litres of product per hectare.
- **STONE AND POME FRUIT TREES:** apply a single treatment before flowering at a dose of 40–50 mL per 100 litres of water (0.04–0.05%), with a spray mixture volume of 700–1,000 L/ha, without exceeding 0.5 litres of product per hectare.
- **TOMATO:** apply only in greenhouses at a rate of 50–75 mL per 100 litres of water (0.05–0.075%). Apply up to 2 applications per season, with a spray mixture volume of 1,000–1,500 L/ha, without exceeding 0.8 litres per hectare of active ingredient per application.

PRE-HARVEST INTERVAL

Aubergine and tomato: 3 days.

Citrus fruits: 30 days.

Other crops: No pre-harvest interval is required.

PRECAUTIONS

To prevent the development of resistance, do not apply this product or any other product containing piriproxyfen more than once per growing season for San José scale and wood lice, and no more than twice for whiteflies.

HAZARD CLASSIFICATION



REGISTRY NUMBER

24,635