

STARLUQ

5 kg 

COMPOSITION

20% w/w Water-soluble Copper (Cu).

USES

A treatment for copper deficiency in all types of crops, particularly suitable for olive, almond, citrus, vine, stone and pome fruit trees, and vegetables.

DOSAGE AND METHOD OF USE

Apply as a standard foliar spray at a rate of 150–250 g per hectolitre of water. Depending on the crop's stage of development and the severity of the deficiency, 2 to 4 spray applications per growing season are recommended. The dose may be increased to 500 g/hL where deemed necessary or where a rapid result is required.

Apply a sufficient amount of spray to ensure that all parts of the plant are completely covered. The amount required will depend on the crop's stage of development and the condition of the foliage.

Fill the tank halfway with water and add the required amount of product, keeping the agitator running until the tank is filled with the remaining water.

It is recommended that you prepare only the amount of mixture you need, and avoid keeping any leftovers for later use.

PRECAUTIONS FOR USE

Avoid carrying out the treatment during the hottest part of the day.

For citrus trees, apply at the start of the growing season and before summer shoots appear.

Take precautions due to the phytotoxicity associated with copper, particularly in humid areas and with certain varieties of fruit trees and other crops.

It can be used in combination with most plant protection products (except calcium polysulphide).

For further information, please contact our Agricultural Technical Department.

HAZARD CLASSIFICATION



16 LUQSA