

ZETAPRID

600 g **COMPOSITION**

Acetamiprid 20% w/w (200 g/kg).
Water-soluble granules (SG).

AUTHORISED USES

CROP	PEST/EFFECT	DOSAGE
Rapeseed	Meligethes, <i>Brassicogethes aeneus</i>	0.2 kg/ha
Apple tree, Peach tree, Pear tree	Aphids, Aphididae	0.25 kg/ha

Apply by mechanical or manual foliar spraying. When applying with a tractor, this must be done with the cab closed. Apply at the doses indicated in the table when signs of infestation appear, following these recommendations:

Rapeseed: Apply a single treatment when the pest is present, before the end of flowering (BBCH 69), using a spray mixture volume of 400–600 L/ha.

Apple and Pear Trees: Except against the species *E. lanigerum* and *A. spiraeicola*. Apply when the pest is present before BBCH 59 and from BBCH 69 onwards, using a spray mixture volume of 1,000 L/ha. Do not apply during flowering. Apply a maximum of once per campaign.

Peach tree: Apply when the pest is present, before growth stage BBCH 59 and from BBCH 69 onwards, using a spray mixture volume of 1,000 L/ha. Do not apply during flowering. Apply no more than once.

PRE-HARVEST INTERVAL

Rapeseed: 42 days.

Apple tree, peach tree, pear tree: 14 days.

PRECAUTIONS

The preparation shall not be used in combination with other products.

It is recommended that a strategy be adopted for the prevention and management of resistance.

In addition, regular assessments should be carried out to monitor any changes in the pests' susceptibility to the product over time.

HAZARD CLASSIFICATION**REGISTRY NUMBER**

ES-00876

